

## RECORDS OF AN INVASIVE DUNG BEETLE SPECIES, *DIGITONTHOPHAGUS GAZELLA* (FABRICIUS, 1787) (COLEOPTERA: SCARABAEIDAE), IN PERU

Jorge Ari NORIEGA<sup>1,2</sup>, Finbarr G. HORGAN<sup>2,3</sup>,  
Trond H. LARSEN<sup>2,4</sup> and Gorky VALENCIA<sup>5</sup>

<sup>1</sup>Laboratorio de Zoología y Ecología Acuática – LAZOEA, Universidad de Los Andes,  
Bogotá, COLOMBIA

<sup>2</sup> Scarabaeinae Research Network - ScarabNet.

<sup>3</sup> International Rice Research Institute, DAPO Box 7777, Metro Manila, THE PHILIPPINES

<sup>4</sup> Department of Ecology and Evolutionary Biology, Princeton University.

<sup>5</sup> Museo de Historia Natural de la Universidad Nacional de San Antonio Abad del Cuzco, PERU.  
E-mail: jnorieg@hotmail.com, f.horgan@cgiar.org; tlarsen@Princeton.EDU; gorkyv@yahoo.com

**Noriega, J. A., F. G. Horgan, T. H. Larsen & G. Valencia.** 2010. Records of an invasive dung beetle species, *Digitonthophagus gazella* (Fabricius, 1787) (Coleoptera: Scarabaeidae), in Peru. *Acta Zool. Mex. (n.s.)*, 26(2): 451-456.

**ABSTRACT.** *Digitonthophagus gazella*, a dung beetle of indoafrikan origin was first introduced to the Americas in the 1970s. Since then it has increased its range through deliberate introductions by national programs and by natural spread in the absence of geographical barriers. The species continues to expand its range as it invades new regions following the anthropogenic clearing of forests. We present the first reports of *D. gazella* in Peru and indicate that the species is now, perhaps, the most widespread dung beetle in tropical and subtropical pastures.

**Key words:** Scarabaeidae, *Digitonthophagus gazella*, dung beetle, invasive species, Peru.

**Noriega, J. A., F. G. Horgan, T. H. Larsen & G. Valencia.** 2010. Registros de una especie invasora de escarabajo coprófago, *Digitonthophagus gazella* (Fabricius, 1787) (Coleoptera: Scarabaeidae), en Perú. *Acta Zool. Mex. (n.s.)*, 26(2): 451-456.

**RESUMEN.** *Digitonthophagus gazella*, un escarabajo coprófago de origen Indoafricano fue inicialmente introducido en el continente Americano en 1970. Desde ese momento su rango de distribución se ha incrementado debido a introducciones deliberadas en nuevas regiones y por procesos naturales de dispersión en ausencia de barreras geográficas. Esta especie ha continuado expandiendo su rango de distribución ayudado por el efecto antrópico producido por la tala de bosques. Se presenta el primer reporte de *D. gazella* para el Perú, indicando que esta especie se constituye en el escarabajo coprófago más ampliamente distribuido en pastizales tropicales y subtropicales.

**Palabras clave:** Scarabaeidae, *Digitonthophagus gazella*, escarabajos coprófagos, especies invasoras, Perú.

---

Recibido: 03/02/2010; aceptado: 20/04/2010.

## INTRODUCTION

*Digitonthophagus gazella* (Fabricius, 1787) (Scarabaeidae: Onthophagini) is a dung beetle species of Indo-african origin that was introduced to Texas (USA) in 1970 for the biological control of bovine dung (Fincher *et al.* 1983). The species has increased its distribution range, helped by anthropic introductions to other countries, colonizing Central and South America and the Caribbean (Huchet 1992, Kohlmann 1994, Gámez *et al.* 1997, Maes *et al.* 1997, Ruiz 2000, Noriega 2002, Noriega *et al.* 2006, Vidaurre *et al.* 2008).

*Digitonthophagus gazella* has been deliberately introduced on a number of occasions to the Neotropics. The initial release in the United States and its subsequent spread through Mexico and Central America has been well documented (Fincher *et al.* 1983, Kohlmann 1994, Maes *et al.* 1997). Its spread through the Caribbean has been less well studied; however, it is now known to occur in Cuba (F. Horgan personal observation), San Andres-Colombia (Noriega 2002), and the Antilles (Huchet 1992). It is perhaps from these regions that it made its way to the Caribbean coast of Colombia (Noriega *et al.* 2006).

The spread of *D. gazella* in South America, other than the Caribbean coast, has been less well documented. The beetle was deliberately introduced into Brazil by the National Center for Beef Cattle Research of the EMBRAPA (Brazilian Enterprise for Research on Agriculture) in 1989, to increase soil productivity in open cattle pastures and as a biological control agent for the horn fly, *Haematobia irritans* (L., 1758) and gastrointestinal helminthes (Oliveira *et al.* 1997, Miranda *et al.* 2000). The first area of liberation was Campo Grande in Mato Grosso do Sul (MS); however, it appears that the species was released over a wide area during the 1990's (i.e., Oliveira *et al.* 1997), and has rapidly built-up numbers and spread from these initial sites. Currently, *D. gazella* features as a prominent species among the recent and extensive documentation of dung beetle communities in Brazilian pastures and grasslands (i.e., Koller *et al.* 1999, Aidar *et al.* 2000, Marchiori *et al.* 2003, Koller *et al.* 2007, Matavelli & Louzada 2008). *Digitonthophagus gazella* is now known to occur in Paraguay, Bolivia and Argentina, presumably through natural spread and has been deliberately introduced into Uruguay and Venezuela (Vidaure *et al.* 2008, Álvarez *et al.* 2009).

In this paper, we present the first records of *D. gazella* in Peru, indicating further expansion of the species' range in South America. Records were compiled from pitfall trapping and direct collections from cow pads, carried out over five years (1999-2003) by the authors in different localities of Madre de Dios and Ucayali.

## RESULTS

**Examined specimens (Fig. 1): PERU.** **Madre de Dios:** 12 (♂ and ♀), Iñapari, 238 m, (12°3' S - 69°24' W), Dec 2002 and May 2003, *F. Horgan leg.*, Universidad Agraria la Molina (UAM). 28 (♂ and ♀), Jorge Chavez near Puerto Maldonado in cattle pastures, 230 m, (12°38'59.4" S - 69°06'24.3" W), Sep 1999, *T. Larsen leg.*, Adrian Forsyth Insect Collection Washington, D.C. (AFIC). 2 ♂, Tahuamanu, near Iñapari, Sep 2002, *F. Horgan leg.*, Entomological Collection of the Museo de Historia Natural de la Universidad Nacional Mayor de San Marcos (MUNMSM). **Ucayali:** 1♂ and 2♀, Ucayali river region near Pucallpa, 160 m, (8°25' S - 74°27' W), Ago 2000, *J.A. Noriega leg.*, Reference Collection Jorge Ari Noriega (CJAN).



**Figure 1.** Localities (●) where *D. gazella* has been registered in Peru. Shaded area indicates Andes mountains.

*Digitonthophagus gazella* appears to be limited to cattle pastures and open savannas at low elevations, it has not been found in the forest interior or in traps placed in nearby forest. *Digitonthophagus gazella* sampled by T. Larsen were captured using two different kinds of bait, human dung and cattle dung,  $2.33 \pm 0.84$  individuals were caught in each 24-hour trap sample based on standard sampling techniques (Larsen & Forsyth 2005). Out of seven dung beetle species sampled in cattle pastures, *D. gazella* was the second most abundant after *Pseudocanthon xanthorus* Blanchard, 1843 (T. Larsen, unpub. data). Similarly, *D. gazella* was recorded in pitfall traps baited with horse and cattle dung in pastures by F. Horgan, with no records from adjacent forest where a similar trapping protocol was conducted. Only, small numbers of *D. gazella* were captured in traps baited with ca. 0.7 kg of dung, possibly because of bait drying in the pastures during the day (baits were set out in the early mornings). Twelve Scarabaeidae species were recorded in the pastures, of which *Gromphas* sp. was the most common.

*Digitonthophagus gazella* has not been sampled above 1000 m in Peru despite extensive sampling by the authors. Assemblages of dung beetles have been sampled thoroughly in grasslands and pastures in other regions of Peru (Huanuco, Tingo Maria, Pasco, Oxapampa, San Ramón, Cusco, Lower and High Urubamba Region); but, *D. gazella* has not been registered in any of these studies (Valencia 2001, Horgan 2005, 2009, Valencia & Concha 2007, Valencia et al. 2009a, b).

## DISCUSSION

The present paper is the first record of the occurrence of *D. gazella* in Peru, notably along Peru's border with Acre-Brazil, a region of extensive cattle production, particularly near Rio Branco in the South of the state (Fig. 1). The records from near Pulcalpa are relatively isolated from Brazilian pastures. Similarly, Matavelli & Louzada (2008) indicate that in Brazil, the species has invaded inter-Amazonian grasslands that are geographically isolated from cattle pastures due to an extensive barrier of rainforest.

The accumulated records on the occurrence of *D. gazella* suggest that the species has invaded new regions in three ways: initially, through deliberate introductions (e.g., the United States, Venezuela, Brazil and Chile); later, by natural dispersal where sites are connected by a series of open pastures (e.g., through Mexico, Central and South America and to some Caribbean islands that are close to the mainland); and finally, through accidental introductions associated with the transportation of livestock in cattle-trucks and river boats (e.g., possibly to inter-Amazonian grasslands, coastal Colombia, Argentina and Ucayali-Peru).

This new report of *D. gazella* in Peruvian territory confirms its status as a highly invasive species; at this time, *D. gazella* is perhaps the most widespread dung beetle in tropical and subtropical pastures. Its ecological success in Neotropical pastures

may be related to its unique functional role as a nocturnal-crepuscular fast tunneller, a functional group that has no native Neotropical members (Horgan 2008), however the possible effects that the species has on the native dung beetle assemblage structure and ecosystem dynamics remains to be studied.

**ACKNOWLEDGEMENTS.** We thank the support of the Laboratory of Zoology and Aquatic Ecology of the Universidad de Los Andes, Natural History Museum of the Universidad Nacional Mayor de San Marcos, Lima and the Natural History Museum of the Universidad Nacional de San Antonio Abad del Cusco, program of Monitoreo de la Biodiversidad en Camisea. The comments and suggestions of two anonymous referees substantially improved the manuscript. We thank ScarabNet for its contribution to the development of better communication mechanisms and aid among researchers.

## LITERATURE CITED

- Aidar, T., W.W. Koller, S.R. Rodrigues, A.M. Corrêa, J.C.C. da Silva, O.S. Balta, J.M. de Oliveira & V.L. de Oliviera.** 2000. Besouros coprófagos (Coleoptera: Scarabaeidae) coletados em Aquidauana, MS, Brasil. *Anais da Sociedade Entomológica do Brasil*, 29(4): 817-820.
- Álvarez, M.C., M.P. Damborsky, M.E. Bar & F.C. Ocampo.** 2009. Registros y distribución de la especie afroasiática *Digitonthophagus gazella* (Coleoptera: Scarabaeidae: Scarabaeinae) en Argentina. *Revista de la Sociedad Entomológica Argentina*, 68(3-4): 373-376.
- Fincher, G.T., T.B. Stewart & J.S. Hunter III.** 1983. The 1981 distribution of *Onthophagus gazella* Fabricius from releases in Texas and *Onthophagus taurus* Schreber from an unknown release in Florida (Coleoptera: Scarabaeidae). *The Coleopterists Bulletin*, 37: 159-163.
- Gámez, J., E. Mora & A. de Ascencao.** 1997. Coleópteros copronecrófilos (Scarabaeidae) en un sistema agropastoril en el sur del Lago de Maracaibo. En: *Resumen XV Congreso Venezolano de Entomología, Trujillo, Venezuela*. 51 pp.
- Horgan, F.G.** 2005. Effects of deforestation on diversity, biomass and function of dung beetles on the eastern slopes of the Peruvian Andes. *Forest Ecology and Management*, 216: 117-133.
- Horgan, F.G.** 2008. Dung beetle assemblages in forests and pastures of El Salvador: A functional comparison. *Biodiversity and Conservation*, 17: 2961-2978.
- Horgan, F.G.** 2009. Invasion and retreat: shifting assemblages of dung beetles amidst changing agricultural landscapes in central Peru. *Biodiversity and Conservation*, 18: 3519-3541.
- Huchet, J.B.** 1992. Un scarabée nouveau pour les Petites Antilles: *Digitonthophagus gazella* (Fabricius) [Col. Scarabaeidae, Coprinae, Onthophagini]. *L'Entomologiste*, 48(6): 297-303.
- Kohlmann, B.** 1994. A preliminary study of the invasion and dispersal of *Digitonthophagus gazella* (Fabricius, 1787) in México (Coleoptera: Scarabaeidae: Scarabaeinae). *Acta Zoológica Mexicana* (n.s.), 61: 35-42.
- Koller, W.W., A. Gomes, S.R. Rodrigues & R.G. De Oliveira.** 1999. Besouros coprófagos (Coleoptera: Scarabaeidae) coletados em Campo Grande, MS. Brasil. *Anais da Sociedade Entomológica do Brasil*, 28(3): 403-412.
- Koller, W.W., A. Gomes, S.R. Rodrigues & P.F. Izique-Goiozo.** 2007. Scarabaeidae e Aphodiidae coprofágos em pastagens cultivadas en area do cerrado sul-mato-grossense. *Revista Brasileira de Zoociências*, 9: 81-93.
- Larsen, T.H. & A. Forsyth.** 2005. Trap spacing and transect design for dung beetle biodiversity studies. *Biotropica*, 37: 322-325.

- Maes, J.M., B.C. Ratcliffe & M.L. Jameson.** 1997. Fauna entomológica de la Reserva Natural Bosawas, Nicaragua. XI. Escarabajos (Coleoptera: Scarabaeidae) nuevos para la fauna de Nicaragua. *Revista Nicaragüense Entomológica*, 39: 41-45.
- Matavelli, R.A. & J.N.C. Louzada.** 2008. Invasão de áreas de savana intra-amazônicas por *Digitonthophagus gazella* (Fabricius, 1787) (Insecta: Coleoptera: Scarabaeidae). *Acta Amazônica*, 38: 153-158.
- Marchiori, C.H., E.R. Caldas & K.G.S. Almeida.** 2003. Succession of Scarabaeidae on bovine dung in Itumbiara, Goiás, Brazil. *Neotropical Entomology*, 32: 173-176.
- Miranda, C.H.B., J.C dos Santos & I. Bianchin.** 2000. The role of *Digitonthophagus gazella* in pasture cleaning and production as a result of burial of cattle dung. *Pasturas Tropicales*, 22(1): 14-18.
- Noriega, J.A.** 2002. First report of the presence of the genus *Digitonthophagus* (Coleoptera: Scarabaeidae) in Colombia. *Caldasia*, 24(1): 213-215.
- Noriega, J.A., C. Solís, I. Quintero, L.G. Perez, H.G. Garcia & D.A. Ospino.** 2006. Registro continental de *Digitonthophagus gazella* (Coleoptera: Scarabaeidae) en Colombia. *Caldasia*, 28(2): 379-381.
- Oliveira, V.B., C.A.D. de Teixeira & F.G. Silva Netto.** 1997. Distribuição do besouro *Onthophagus gazella* (Scarabaeidae) para o CB da mosca-dos-chifres (*Haematobia irritans* - Muscidae) em Rondônia. In: *Resumos Sociedade Brasileira de Entomologia/EMBRAPA-CNPMF. Congresso Brasiliero de Entomologia 16.*, Salvador.
- Ruiz, M.A.** 2000. *Levantamento populacional de besouros coprofagos (Coleoptera Scarabaeidae) no Estado de Amambay, Repùblica do Paraguai.* Dissertação de Mestrado em Entomologia, Escola Superior de Agricultura “Luis de Queiroz”, Universidade de São Paulo, Piracicaba, Brasil. 80 pp.
- Valencia, G.** 2001. Diversity and trophic relationships of dung beetles of the well sites from Lower Urubamba Region, Perú. In: F. Dallmeier, A. Alonso & P. Campbell. (Eds.). *Urubamba: The Biodiversity of a Peruvian Rainforest.* SI/MAB series # 7. Smithsonian Institution, Washington, D.C.
- Valencia, G. & R.D. Concha.** 2007. *Diversidad de artrópodos en San Martín y Las Malvinas, con énfasis en Coleoptera Scarabaeoidea.* (PMB. Época de lluvias). Informe, ERM – Perú. 53 pp.
- Valencia, G., R.D. Concha & M. Cardenas.** 2009a. *Diversidad de artrópodos en el flowline de la Planta de Gas de las Malvinas (Kp 50 Y 84).* PMB downstream septiembre del 2008. Informe, ERM – Perú. 25 pp.
- Valencia, G., R.D. Concha & M. Cardenas.** 2009b. *Diversidad de artrópodos en Mipaya y San Martín 3.* PMB upstream febrero 2009. Informe, ERM – Perú. 50 pp.
- Vidaurre, T., J.A. Noriega & M.J. Ledezma.** 2008. First report on the distribution of *Digitonthophagus gazella* (Fabricius, 1787) (Coleoptera: Scarabaeidae) in Bolivia. *Acta Zoológica Mexicana*, 24(3): 217-220.